



HAL
open science

Efficacy of artesunate–amodiaquine in the treatment of falciparum uncomplicated malaria in Madagascar

Oméga Raobela, Valérie Andriantsoanirina, David Gael Rajaonera, Tovonahary Angelo Rakotomanga, Stéphane Rabearimanana, Fanomezantsoa Ralinoro, Didier Ménard, Arsène Ratsimbasa

► To cite this version:

Oméga Raobela, Valérie Andriantsoanirina, David Gael Rajaonera, Tovonahary Angelo Rakotomanga, Stéphane Rabearimanana, et al.. Efficacy of artesunate–amodiaquine in the treatment of falciparum uncomplicated malaria in Madagascar. *Malaria Journal*, 2018, 17, pp.284. 10.1186/s12936-018-2440-0 . hal-02558706

HAL Id: hal-02558706

<https://hal.science/hal-02558706>

Submitted on 6 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

RESEARCH

Open Access



Efficacy of artesunate–amodiaquine in the treatment of falciparum uncomplicated malaria in Madagascar

Oméga Raobela^{1†}, Valérie Andriantsoanirina^{1*†}, David Gael Rajaonera¹, Tovonahary Angelo Rakotomanga^{1,2}, Stéphane Rabearimanana^{1,2}, Fanomezantsoa Ralinoro^{1,2}, Didier Ménard³ and Arsène Ratsimbaoa^{1,4}

Abstract

Background: Since 2006, the artemisinin-based combination therapy (ACT) are recommended to treat uncomplicated malaria including non *Plasmodium falciparum* malaria in Madagascar. Artesunate–amodiaquine (ASAQ) and artemether–lumefantrine are the first- and second-line treatment in uncomplicated falciparum malaria, respectively. No clinical drug efficacy study has been published since 2009 to assess the efficacy of these two artemisinin-based combinations in Madagascar, although the incidence of malaria cases has increased from 2010 to 2016. In this context, new data about the efficacy of the drug combinations currently used to treat malaria are needed.

Methods: Therapeutic efficacy studies evaluating the efficacy of ASAQ were conducted in 2012, 2013 and 2016 among falciparum malaria-infected patients aged between 6 months and 56 years, in health centres in 6 sites representing different epidemiological patterns. The 2009 World Health Organization protocol for monitoring anti-malarial drug efficacy was followed.

Results: A total of 348 enrolled patients met the inclusion criteria including 108 patients in 2012 (n = 64 for Matanga, n = 44 for Ampasipotsoy), 123 patients in 2013 (n = 63 for Ankazomborona, n = 60 for Anjoma Ramartina) and 117 patients in 2016 (n = 67 for Tsaratanana, n = 50 for Antanimbary). The overall cumulative PCR-corrected day 28 cure rate was 99.70% (95% IC 98.30–99.95). No significant difference in cure rates was observed overtime: 99.02% (95% IC 94.65–99.83) in 2012; 100% (95% IC 96.8–100) in 2013 and 100% (95% IC 96.65–100) in 2016.

Conclusion: The ASAQ combination remains highly effective for the treatment of uncomplicated falciparum malaria in Madagascar.

Keywords: Malaria, *Plasmodium falciparum*, Therapeutic efficacy survey, Artesunate–amodiaquine, Madagascar

Background

Malaria remains an important health problem in Madagascar mainly in children under 5 years of age as mentioned in the national strategy plan for malaria control in Madagascar 2013–2017 [1]. Over the past decade, the burden of malaria has fluctuated over time partly due to successes and failures of anti-malarial policy [1]. The

artemisinin-based combination therapy (ACT) are recommended in Madagascar since 2006 to treat uncomplicated malaria including non-falciparum malaria. The National Malaria Control Programme (NMCP) replaced chloroquine with artesunate–amodiaquine (ASAQ) combination as first-line drug for treating uncomplicated falciparum malaria, and artemether–lumefantrine as an alternative treatment. This change was based on a study, including clinical and in vitro data that reported the complete efficacy of the ASAQ combination in Madagascar [2, 3]. An additional study performed 1 year later, which was part of a multicentre trial conducted in 2006 in 5 African sites, reported similar high levels of cure rates

*Correspondence: landyvalerie@gmail.com

†Oméga Raobela and Valérie Andriantsoanirina contributed equally to this work

¹ National Malaria Control Programme of Madagascar, Androhibe, Antananarivo, Madagascar

Full list of author information is available at the end of the article



to ASAQ in Tsiroanomandidy [4]. Since then, no data about the efficacy of ASAQ or alternative anti-malarials in Madagascar have been reported.

The incidence of malaria cases has increased in Madagascar (9.83 cases for 1000 inhabitants in 2010 to 19.52 in 2016) (NMCP data, pers. comm.). In this context, it is of utmost importance to acquire new data from recent efficacy studies. This study presents the results of studies conducted from 2012 to 2016 to assess the efficacy of the ASAQ combination for the treatment of uncomplicated malaria cases in 6 sites covering three epidemiological patterns in Madagascar.

Methods

Study areas

Clinical ASAQ efficacy studies were carried out in 6 sites belonging to 3 epidemiological strata, from May to September 2012 in Matanga (Vangaindrano) and Ampasimpotsy (Tsiroanomandidy), from March to May 2013 in Ankazomborona (Marovoay) and Anjoma Ramartina (Mandoto), and from March to June 2016 in Tsaratanana (Ifanadiana) and Antanimbary (Maevatanana) (Fig. 1).

Matanga and Tsaratanana are located in eastern Madagascar (equatorial stratum, perennial and endemic area), Antanimbary (Maevatanana) and Ankazomborona (Marovoay) in western Madagascar (tropical stratum, seasonal and endemic area) and Ampasimpotsy (Tsiroanomandidy) and Anjoma Ramartina (Mandoto) in the fringe highlands of Madagascar (fringe highlands stratum, seasonal and low endemic area).

Patient recruitment

Clinical ASAQ efficacy studies were conducted according to 2009 World Health Organization protocol for monitoring anti-malarial drug efficacy [5]. Febrile patients or patients with fever history seeking anti-malarial treatment in health centres were screened for malaria by rapid diagnostic test (SD Boline Ag P.f/Pan, Standard Diagnostics INC, Korea). Positive cases for *Plasmodium falciparum* were enrolled if they met inclusion criteria and gave their written informed consent. Inclusion criteria were: (i) age > 6 months; (ii) axillary temperature ≥ 37.5 °C or history of fever in the 24 h preceding consultation; (iii) *P. falciparum* mono-infection with parasitaemia between 1000 and 200,000 asexual parasites per μ l; and, (iv) absence of signs of severe malaria. Children were recruited after parental consent.

Finger-prick capillary blood samples were collected to prepare thin and thick blood smears for malaria microscopy examination, haemoglobin level determination (HemoCue HB 201, HemoCue AB, Angelholm, Sweden) and dried blood spot (DBS) for molecular biology investigations.

Artesunate–amodiaquine Winthrop® (Sanofi, France) was administered daily for 3 days at the dose recommended by the manufacturer on a mg per weight basis (4.5–8 kg: 25 mg/67.5 mg tablets; 9–17.9 kg: 50 mg/135 mg tablets; 18–35 kg: 100 mg/270 mg tablets; > 36 kg: 100 mg/270 mg tablets). Participants were observed by the medical team for 30 min after treatment to monitor for vomiting or other adverse events; those who vomited were administered a second dose and observed for an additional 30 min. Patients who vomited both doses were excluded from the study and referred to the hospital for parenteral treatment.

Patient follow-up

Enrolled patients were followed daily (days 1, 2, 3) and weekly (days 7, 14, 21, and 28). At each visit, patients were clinically examined by a physician and clinical parameters were recorded on a Case Record Form (CRF). Parasite densities estimated by microscopic examination were checked at each visit. All blood films were read by 2 qualified microscopists, and by a third independent microscopist if discordance was > 20%. Dried blood spot obtained at day 0 and day failure were used for PCR genotyping. Haemoglobin concentrations were assessed on day 0 and day 28.

DNA extraction and molecular investigations

Parasite DNA extracted from dried blood spot (DBS) with QIAamp DNA blood kit according to manufacturer's instruction, was amplified for confirming *Plasmodium* species with slight modifications [6] and for distinguishing recrudescence and re-infection by using *m*sp1 and *m*sp2 polymorphisms [7] in case of recrudescence.

Data analysis

Data were entered into the standard pre-programmed Excel worksheet provided by the Global Malaria Programme WHO for per-protocol analysis. Per protocol analysis was used to assess treatment outcomes at day 28 based on the WHO 2009 criteria: ACPR (adequate clinical and parasitological response), ETF (early treatment failure), LCF (late clinical failure), and LPF (late parasitological failure).

Results

A total of 348 patients met the inclusion criteria and were enrolled for the analyses including 108 patients in 2012 (n = 64 for Matanga, n = 44 for Ampasimpotsy), 123 patients in 2013 (n = 63 for Ankazomborona, n = 60 for Anjoma Ramartina) and 117 patients in 2016 (n = 67 for Tsaratanana, n = 50 for Antanimbary). Their baseline characteristics are shown in Table 1. Ages ranged from 6 months to 56 years (median 8 years).




Table 1 Baseline clinical and parasitological characteristics of enrolled patients in the 6 sites, Madagascar, 2012–2016, ASAQ

Epidemiological strata	Eastern Madagascar (equatorial stratum)		Western Madagascar (tropical stratum)		Fringe highlands of Madagascar (fringe highlands stratum)	
	Matanga	Tsatanana	Ankazomborona	Antanimbary	Ampasipotsy	Anjoma Ramartina
Site	Matanga	Tsatanana	Ankazomborona	Antanimbary	Ampasipotsy	Anjoma Ramartina
District	Vangaindrano	Ifanadiana	Marovoay	Maevatanana	Tsiroanomandidy	Mandoto
Year	2012	2016	2013	2016	2012	2013
n	64	67	63	50	44	60
Age years, median (range)	2.17 (1–34)	6 (1–14)	9 (1–54)	11 (1–56)	174 (2–56)	10 (2–45)
Gender male, n (%)	35 (54.69)	24 (35.82)	29 (46.03)	28 (35.82)	27 (61.36)	30 (50)
Axillary T°, mean (SD)	38.05 (1.38)	38.42 (1.02)	37.83 (1.46)	38.38 (1.29)	37.91 (1.14)	37.96 (1.31)
Parasite density/ μ l	20,803.67	28,089.02	21,517.67	22,301.17	7315.71	10,704.84
Parasite density (geometric mean, range)	1005–199,800	1561–185,103	1448–198,400	1168–151,844	1056–85,000	1009–149,747
Age group, n (%)						
Under 5	55 (85.94)	25 (37.35)	13 (20.63)	7 (14)	3 (6.82)	13 (21.70)
5–15	8 (12.5)	42 (62.65)	35 (55.56)	28 (56)	19 (43.18)	30 (50)
Adult	1 (1.56)	0 (0)	15 (23.81)	15 (30)	22 (50)	17 (28.30)
Haemoglobin g/dl (mean, SD)	ND	10.24 (1.90)	11.25 (1.90)	10.76 (1.92)	ND	10.43 (2.21)
Hb < 10 g/dl, n (%)	ND	29 (43.3)	14 (22.23)	16 (32)	ND	21 (35)

ND not done, SD standard deviation

The geometric mean of the parasite density at day 0 was 17,502 parasites/ μ l of blood (95% CI 13,000–22,003, range 1005–199,800). “The parasite density mean varied according to sites, from 7315 parasites/ μ l of blood in Ampasipotsy to 28,089 parasites/ μ l of blood in Tsatanana”.

The combination was well tolerated. Few adverse events in gastrointestinal transit (vomiting, nausea, abdominal pain, anorexia), in the central nervous systems (headache, dizziness, drowsiness) and in musculoskeletal system were observed in 11.78% (41/348) patients, 2.58% (9/348) and 2.58% (9/348), respectively. The median haemoglobin concentration, among patients with available data (196/358), significantly increased from 10.60 g/dl on day 0 to 11.50 g/dl on day 28 ($P < 0.00001$).

The proportion of patients still harbouring parasites over the first 3 days were 42.65% (148/347) at day 1, 2.31% (8/346) at day 2 and 0% (0/340) at day 3, respectively.

The overall efficacy of ASAQ per protocol population analysis and after PCR correction was 99.70% (95% IC 98.30–99.95). According to years, PCR-corrected day 28 cure rates were 99.02% (95% IC 94.65–99.83) in 2012; 100% (95% IC 96.8–100) in 2013 and 100% (95% IC 96.65–100) in 2016. There was no case of ETF; 2 cases of LPF and one case of LCF were observed. The first patient classified as LPF was a 17-month old boy with a parasitaemia of 18,613 asexual parasites/ μ l at day 0 and 150,000 asexual parasites/ μ l at day 28. This patient was a true

recrudescence as determined by *msp1/msp2* genotyping. The second patient classified as LPF was a 17-month old girl with a parasitaemia of 9900 asexual parasites/ μ l at day 0 and 29,232 asexual parasites/ μ l at day 28. The patient presenting a LCF, a 10-year old male, had a parasitaemia of 5811 asexual parasites/ μ l on day 0 and 9400 asexual parasites/ μ l on day 28 with 38.9 °C of axillary temperature. The latter two patients were classified as reinfection by *msp1/msp2* genotyping and were excluded from the analysis (Table 2).

Discussion

Amodiaquine, in combination with artesunate, is widely used in sub-Saharan African countries, including Madagascar. The aim of this study was to provide updated data about the efficacy of ASAQ combination in Madagascar. Twelve years after its adoption as first-line treatment for acute uncomplicated falciparum malaria, ASAQ remains highly efficacious in the 6 geographical sites, covering various eco-epidemiological facies of the island, with different levels of malaria transmission. PCR-corrected cured rates estimated at day 28 were largely > 90%, above the WHO threshold for changing malaria treatment policy [5]. In addition, all patients treated with ASAQ cleared their parasites before day 3, indicating the absence of delayed parasite clearance, a marker for suspected partial resistance to artemisinin [8].

Table 2 Treatment outcomes (per protocol analysis) by epidemiological strata, Madagascar 2012–2016, ASAQ

Epidemiological strata	Eastern Madagascar (equatorial stratum)			
Site	Matanga		Tsaratana	
District	Vangaindrano		Ifanadiana	
Year	2012		2016	
Number of patients, n	64		67	
Clinical outcomes	Unadjusted-PCR	PCR-corrected	Unadjusted-PCR	PCR-corrected
ETF, n	0	0	0	0
LCT, n	0	0	0	0
LPF, n	2	1	0	0
ACPR, n (%)	62 (96.9)	62 (98.4)	65 (100)	65 (100)
CI 95% (%)	89.3–99.1	91.5–99.7	94.4–100	94.4–100
Withdrawal, n	0	1	2 ^a	2 ^a
Lost to follow-up, n	0	0	0	0
Epidemiological strata	Fringe highlands of Madagascar (fringe highlands stratum)			
Site	Ampasipotry		Anjoma Ramartina	
District	Tsiroanomandidy		Mandoto	
Year	2012		2013	
Number of patients, n	44		60	
Clinical outcomes	Unadjusted-PCR	PCR corrected	Unadjusted-PCR	PCR corrected
ETF	0	0	0	0
LCT	1	0	0	0
LPF	0	0	0	0
ACPR	39 (97.5)	39 (100)	53 (100)	53 (100)
CI 95% (%)	87.1–99.6	91–100	93.2–100	93.2–100
Withdrawal, n	0	1	0	0
Lost to follow-up, n	4	4	7	7
Epidemiological strata	Western Madagascar (tropical stratum)			
Site	Ankazomborona		Antanimbary	
District	Marovoay		Maevatanana	
Year	2013		2016	
Number of patients, n	63		50	
Clinical outcomes	Unadjusted-PCR	PCR corrected	Unadjusted-PCR	PCR corrected
ETF	0	0	0	0
LCT	0	0	0	0
LPF	0	0	0	0
ACPR	63 (100)	63 (100)	46 (100)	46 (100)
CI 95% (%)	94.3–100	94.3–100	92.3–100	92.3–100
Withdrawal, n	0	0	0	0
Lost to follow-up, n	0	0	4	4

ETF early therapeutic failure, LCF late clinical failure, LPF late parasitological failure, ACPR adequate clinical and parasitological response

^a Withdrawal of consent

The proportion of cured patients observed in this study was not different from that reported in the two studies performed a decade earlier, suggesting the ASAQ retained its efficacy so far. These results are in line with those of comparable studies performed

recently in neighbouring countries, such as Mozambique or Kenya [9, 10]. The results obtained in this study are also in line with those obtained in several African countries where ASAQ therapeutic failures are very rarely reported [11–18]. When documented, those

therapeutic failures were not associated with the same *k13* mutations that were reported in *P. falciparum* isolates obtained during therapeutic failures in Cambodia and other Southeast Asian countries [19–22].

The treatment by ASAQ was well tolerated and no severe adverse event was reported among the participants in the 6 sites.

The efficacy of ASAQ, demonstrated in the present study, is necessary but not sufficient to contribute to the elimination of malaria in Madagascar. Stock out issues are recurrent on the island, emphasizing that sustainability of distribution of anti-malarials is crucial.

Conclusion

ASAQ remains highly efficacious in the treatment of uncomplicated falciparum malaria in Madagascar. Thus, according to this study, the increased incidence of falciparum malaria cases observed recently seems not related to clinical treatment failure of first-line treatment promoted since 2006. Further investigations are required to determine the reasons for the recent increase in incidence of falciparum malaria.

Authors' contributions

OR and AR conceived, designed, coordinated the study, supervised the enrolment and follow up of the patients. VA, DGR and TAR analysed the data and map. SR and FR performed the molecular assay. VA, AR and DM composed the manuscript. All authors read and approved the final manuscript.

Author details

¹ National Malaria Control Programme of Madagascar, Androhibe, Antananarivo, Madagascar. ² Faculty of Sciences, University of Antananarivo, Antananarivo, Madagascar. ³ Département Parasites et Insectes Vecteurs, Institut Pasteur Paris, Paris, France. ⁴ Faculty of Medicine, University of Antananarivo, Antananarivo, Madagascar.

Acknowledgements

We thank all study participants, their guardians and the parents of children for their participation in these studies, the health care staff in each of the six health facilities and study team members: Michel Abraham Marolahy, Tiavina Rakotomanjaka, Brunette Razanadrazanina, Noeline Rasoarilalao, Dr. Marie-Ange Rason, Jaona Rahajarimanitriainaina Claude L.G., Yvon Ralaiseheno.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

All clinical drug efficacy studies were conducted in accordance with the Helsinki Declaration and the national laws and regulations in force. The protocol was reviewed and approved by the Malagasy National Ethics Committee (No. 083/MSANP/CE/11-2012). A written informed consent was obtained from patients or the parents/guardians of all children before inclusion in the study.

Funding

This work was supported by The Global Fund: NSA2 (MDG-M-PSI 635) and NMF1 (MDG-M-PSI 1044).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 12 April 2018 Accepted: 1 August 2018

Published online: 06 August 2018

References

- PNLP. Plan stratégique de lutte contre le paludisme à Madagascar, 2013–2017. Antananarivo; 2013.
- Menard D, Ratsimbao A, Randrianarivojosia M, Rabarijaona LP, Rahrimalala L, Domarle O, et al. Assessment of the efficacy of antimalarial drugs recommended by the National Malaria Control Programme in Madagascar: up-dated baseline data from randomized and multi-site clinical trials. *Malar J*. 2008;7:55.
- Andriantsoanirina V, Ratsimbao A, Bouchier C, Jahevitra M, Rabearimanana S, Radrianjafy R, et al. *Plasmodium falciparum* drug resistance in Madagascar: facing the spread of unusual pfdhfr and pfmdr-1 haplotypes and the decrease of dihydroartemisinin susceptibility. *Antimicrob Agents Chemother*. 2009;53:4588–97.
- Ndiaye JL, Randrianarivojosia M, Sagara I, Brasseur P, Ndiaye I, Faye B, et al. Randomized, multicentre assessment of the efficacy and safety of ASAQ—a fixed-dose artesunate–amodiaquine combination therapy in the treatment of uncomplicated *Plasmodium falciparum* malaria. *Malar J*. 2009;8:125.
- WHO. Methods for surveillance of antimalarial drug efficacy. Geneva: World Health Organization; 2009.
- Snounou G. Detection and identification of the four malaria parasite species infecting humans by PCR amplification. *Methods Mol Biol*. 1996;50:263–91.
- Snounou G, Zhu X, Siripoon N, Jarra W, Thaitong S, Brown KN, et al. Biased distribution of msp1 and msp2 allelic variants in *Plasmodium falciparum* populations in Thailand. *Trans R Soc Trop Med Hyg*. 1999;93:369–74.
- WHO. Global Malaria Programme. Status report on artemisinin resistance. Geneva: World Health Organization; 2014. <http://www.who.int/malaria/publications/atoz/status-rep-artemisinin-resistance-sep2014.pdf>.
- Thwing JI, Odero CO, Odhiambo FO, Otieno KO, Kariuki S, Ord R, et al. In vivo efficacy of amodiaquine-artesunate in children with uncomplicated *Plasmodium falciparum* malaria in western Kenya. *Trop Med Int Health*. 2009;14:294–300.
- Nhama A, Bassat Q, Enosse S, Nhacolo A, Mutemba R, Carvalho E, et al. In vivo efficacy of artemether–lumefantrine and artesunate–amodiaquine for the treatment of uncomplicated falciparum malaria in children: a multisite, open-label, two-cohort, clinical trial in Mozambique. *Malar J*. 2014;13:309.
- Dorkenoo AM, Yehadj D, Agbo YM, Layibo Y, Agbeko F, Adjelloh P, et al. Therapeutic efficacy trial of artemisinin-based combination therapy for the treatment of uncomplicated malaria and investigation of mutations in k13 propeller domain in Togo, 2012–2013. *Malar J*. 2016;15:331.
- Mekonnen SK, Medhin G, Berhe N, Clouse RM, Aseffa A. Efficacy of artemether–lumefantrine therapy for the treatment of uncomplicated *Plasmodium falciparum* malaria in southwestern Ethiopia. *Malar J*. 2015;14:317.
- Charle P, Berzosa P, de Lucio A, Raso J, Nseng Nchama G, Benito A. Artesunate/amodiaquine malaria treatment for Equatorial Guinea (Central Africa). *Am J Trop Med Hyg*. 2013;88:1087–92.
- Ouldabdallahi M, Alew I, Salem MS, Dit Dialaw Ba M, Boukhary AO, Khairy ML, et al. Efficacy of artesunate–amodiaquine for the treatment of acute uncomplicated falciparum malaria in southern Mauritania. *Malar J*. 2014;13:496.
- Tahar R, Almelli T, Debue C, Foumane Ngane V, Djaman Allico J, Whengang Youdom S, et al. Randomized trial of artesunate–amodiaquine, atovaquone–proguanil, and artesunate–atovaquone–proguanil for the treatment of uncomplicated falciparum malaria in children. *J Infect Dis*. 2014;210:1962–71.
- Zwang J, Olliaro P, Barennes H, Bonnet M, Brasseur P, Bukirwa H, et al. Efficacy of artesunate–amodiaquine for treating uncomplicated falciparum malaria in sub-Saharan Africa: a multi-centre analysis. *Malar J*. 2009;8:203.
- Espie E, Lima A, Atua B, Dhorda M, Flevaud L, Sompwe EM, et al. Efficacy of fixed-dose combination artesunate–amodiaquine versus artemether–lumefantrine for uncomplicated childhood *Plasmodium falciparum*

- malaria in Democratic Republic of Congo: a randomized non-inferiority trial. *Malar J*. 2012;11:174.
18. Yeka A, Lameyre V, Afizi K, Fredrick M, Lukwago R, Kamya MR, et al. Efficacy and safety of fixed-dose artesunate–amodiaquine vs. artemether–lumefantrine for repeated treatment of uncomplicated malaria in Ugandan children. *PLoS One*. 2014;9:e113311.
 19. Menard D, Khim N, Beghain J, Adegnikaa AA, Shafiu-Alam M, Amodu O, et al. A worldwide map of *Plasmodium falciparum* K13-propeller polymorphisms. *N Engl J Med*. 2016;374:2453–64.
 20. Takala-Harrison S, Jacob CG, Arze C, Cummings MP, Silva JC, Don-dorp AM, et al. Independent emergence of artemisinin resistance mutations among *Plasmodium falciparum* in southeast Asia. *J Infect Dis*. 2015;211:670–9.
 21. Myint MK, Rasmussen C, Thi A, Bustos D, Ringwald P, Lin K. Therapeutic efficacy and artemisinin resistance in northern Myanmar: evidence from in vivo and molecular marker studies. *Malar J*. 2017;16:143.
 22. Thanh NX, Trung TN, Phong NC, Quang HH, Dai B, Shanks GD, et al. The efficacy and tolerability of artemisinin-piperaquine (Artequick(R)) versus artesunate–amodiaquine (Coarsucam) for the treatment of uncomplicated *Plasmodium falciparum* malaria in south-central Vietnam. *Malar J*. 2012;11:217.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

